

DIRECT Distances to Nearby Galaxies Using Detached Eclipsing Binaries and Cepheids. V. Variables in the Field M31F¹

B. J. Mochejska and J. Kaluzny

Warsaw University Observatory, Al. Ujazdowskie 4, PL-00-478 Warszawa, Poland

e-mail: mochejsk@sirius.astrow.edu.pl, jka@sirius.astrow.edu.pl

K. Z. Stanek², M. Krockenberger and D. D. Sasselov

Harvard-Smithsonian Center for Astrophysics, 60 Garden St., Cambridge, MA 02138

e-mail: kstanek@cfa.harvard.edu, krocken@cfa.harvard.edu,
sasselov@cfa.harvard.edu

ABSTRACT

We undertook a long term project, DIRECT, to obtain the direct distances to two important galaxies in the cosmological distance ladder – M31 and M33 – using detached eclipsing binaries (DEBs) and Cepheids. While rare and difficult to detect, DEBs provide us with the potential to determine these distances with an accuracy better than 5%. The extensive photometry obtained in order to detect DEBs provides us with good light curves for the Cepheid variables. These are essential to the parallel project to derive direct Baade-Wesselink distances to Cepheids in M31 and M33. For both Cepheids and eclipsing binaries, the distance estimates will be free of any intermediate steps.

As a first step in the DIRECT project, between September 1996 and October 1997 we obtained 95 full/partial nights on the F. L. Whipple Observatory 1.2 m telescope and 36 full nights on the Michigan-Dartmouth-MIT 1.3 m telescope to search for DEBs and new Cepheids in the M31 and M33 galaxies. In this paper, fifth in the series, we present the catalog of variable stars found in the field M31F $[(\alpha, \delta) = (10^{\circ}10, 40^{\circ}72), J2000.0]$. We have found 64 variable stars: 4 eclipsing binaries, 52 Cepheids and 8 other periodic, possible long period or non-periodic variables. The catalog of variables, as well as their photometry and finding charts, is available via **anonymous ftp** and the **World Wide Web**. The complete set of the CCD frames is available upon request.

Subject headings: binaries: eclipsing — Cepheids — distance scale — galaxies: individual (M31) — stars: variables: other

¹Based on the observations collected at the F. L. Whipple Observatory (FLWO) 1.2 m telescope and at the Michigan-Dartmouth-MIT (MDM) 1.3 m telescope

²On leave from N. Copernicus Astronomical Center, Bartycka 18, Warszawa PL-00-716, Poland

1. Introduction

Starting in 1996 we undertook a long term project, DIRECT (as in “direct distances”), to obtain the distances to two important galaxies in the cosmological distance ladder – M31 and M33 – using detached eclipsing binaries (DEBs) and Cepheids. These two nearby galaxies are stepping stones to most of our current effort to understand the evolving universe at large scales. First, they are essential to the calibration of the extragalactic distance scale (Jacoby et al. 1992; Tonry et al. 1997). Second, they constrain population synthesis models for early galaxy formation and evolution and provide the stellar luminosity calibration. There is one simple requirement for all this—accurate distances.

DEBs have the potential to establish distances to M31 and M33 with an unprecedented accuracy of better than 5% and possibly to better than 1%. These distances are now known to no better than 10-15%, as there are discrepancies of 0.2 – 0.3 mag between various distance indicators (e.g. Huterer, Sasselov & Schechter 1995; Holland 1998; Stanek & Garnavich 1998). Detached eclipsing binaries (for reviews see Andersen 1991; Paczyński 1997) offer a single step distance determination to nearby galaxies and may therefore provide an accurate zero point calibration—a major step towards very accurate determination of the Hubble constant, presently an important but daunting problem for astrophysicists. A DEB system was recently used by Guinan et al. (1998) and Udalski et al. (1998) to obtain an accurate distance estimate to the Large Magellanic Cloud.

The detached eclipsing binaries have yet to be used (Huterer et al. 1995; Hilditch 1996) as distance indicators to M31 and M33. According to Hilditch (1996), there were about 60 eclipsing binaries of all kinds known in M31 (Gaposchkin 1962; Baade & Swope 1963, 1965) and only *one* in M33 (Hubble 1929), none of them observed with CCDs. Only now does the availability of large-format CCD detectors and inexpensive CPUs make it possible to organize a massive search for periodic variables, which will produce a handful of good DEB candidates. These can then be spectroscopically followed-up with the powerful new 6.5-10 meter telescopes.

The study of Cepheids in M31 and M33 has a venerable history (Hubble 1926, 1929; Gaposchkin 1962; Baade & Swope 1963, 1965). Freedman & Madore (1990) and Freedman, Wilson & Madore (1991) obtained multi-band CCD photometry of some of the already known Cepheids, to build period-luminosity relations in M31 and M33, respectively. However, both the sparse photometry and the small samples (11 Cepheids in M33 and 38 Cepheids in M31) do not provide a good basis for obtaining direct Baade-Wesselink distances (see, e.g., Krockenberger, Sasselov & Noyes 1997) to Cepheids—the need for new digital photometry has been long overdue. Recently, Magnier et al. (1997) surveyed large portions of M31, which have previously been ignored, and found some 130 new Cepheid variable candidates. Their light curves are, however, rather sparsely sampled and in the *V*-band only.

In Kaluzny et al. (1998, 1999, hereafter: Papers I and IV) and Stanek et al. (1998, 1999, hereafter: Papers II and III), the first four papers of the series, we presented the catalogs of variable stars found in four fields in M31, called M31B, M31A, M31C and M31D. Here we present

the catalog of variables from the field M31F. In Sec.2 we discuss the selection of the fields in M31 and the observations. In Sec.3 we describe the data reduction and calibration. In Sec.4 we discuss briefly the automatic selection we used for finding the variable stars. In Sec.5 we discuss the classification of the variables. In Sec.6 we present the catalog of variable stars, followed by a brief discussion of the results in Sec.7.

2. Fields selection and observations

M31 was primarily observed in 1996 with the 1.3 m McGraw-Hill Telescope at the Michigan-Dartmouth-MIT (MDM) Observatory. We used the front-illuminated, Loral 2048² CCD “Wilbur” (Metzger, Tonry & Luppino 1993), which at the $f/7.5$ station of the 1.3 m telescope has a pixel scale of $0.32 \text{ arcsec pixel}^{-1}$ and field of view of roughly 11 arcmin . We used Kitt Peak Johnson-Cousins *BVI* filters. Data for M31 were also obtained, mostly in 1997, with the 1.2 m telescope at the F. L. Whipple Observatory (FLWO), where we used “AndyCam” (Szentgyorgyi et al. 1999), with a thinned, back-illuminated, AR coated Loral 2048² pixel CCD. The pixel scale happens to be essentially the same as at the MDM 1.3 m telescope. We used standard Johnson-Cousins *BVI* filters.

Fields in M31 were selected using the MIT photometric survey of M31 by Magnier et al. (1992) and Haiman et al. (1994) (see Paper I, Fig.1). We selected six $11' \times 11'$ fields, M31A–F, four of them (A–D) concentrated on the rich spiral arm in the northeast part of M31, one (E) coinciding with the region of M31 searched for microlensing by Crotts & Tomaney (1996), and one (F) containing the giant star formation region known as NGC206 (observed by Baade & Swope 1965). Fields A–C were observed during September and October 1996 five to eight times per night in the *V* band, resulting in total of 110–160 *V* exposures per field. Fields D–F were observed once a night in the *V*-band. Some exposures in *B* and *I* were also taken. M31 was also observed, in 1996 and 1997, at the FLWO 1.2 m telescope, whose main target was M33.

In this paper we present the results for the M31F field. We obtained for this field useful data during 29 nights at the MDM, collecting a total of $28 \times 900 \text{ sec}$ exposures in *V* and $2 \times 600 \text{ sec}$ exposures in *I*. We also obtained for this field useful data during 22 nights at the FLWO, in 1996 and 1997, collecting a total of $80 \times 900 \text{ sec}$ exposures in *V*, $67 \times 600 \text{ sec}$ exposures in *I* and $7 \times 1200 \text{ sec}$ exposures of *B*.³

³The complete list of exposures for this field and related data files are available through **anonymous ftp** on cfa-ftp.harvard.edu, in `pub/kstanek/DIRECT` directory. Please retrieve the `README` file for instructions. Additional information on the DIRECT project is available through the WWW at <http://cfa-www.harvard.edu/~kstanek/DIRECT/>.

3. Data reduction, calibration and astrometry

The details of the reduction procedure were given in Paper I. Preliminary processing of the CCD frames was done with the standard routines in the IRAF-CCDPROC package.⁴ Stellar profile photometry was extracted using the *Daophot/Allstar* package (Stetson 1987, 1992). We selected a “template” frame for each filter using a single frame of particularly good quality. These template images were reduced in a standard way (Paper I). Other images were reduced using *Allstar* in the fixed-position-mode using as an input the transformed object list from the template frames. For each frame the list of instrumental photometry derived for a given frame was transformed to the common instrumental system of the appropriate “template” image. Photometry obtained for the B , V and I filters was combined into separate data bases. M31F I -band images obtained at the MDM were reduced using FLWO “templates”. Two templates were used in the case of V -band images. An MDM template was used to fix the positions of the stars. The V photometry was transformed to the instrumental system of an FLWO template.

The photometric VI calibration of the MDM data was discussed in Paper I. In addition, for the field M31F on the night of 1997 October 9/10 we have obtained independent BVI calibration with the FLWO 1.2 m telescope. There was an offset of -0.020 mag in V and 0.057 mag in $V - I$ between the FLWO and the MDM calibration. The V offset is well within our estimate of the total 0.05 mag systematic error discussed in Paper I, and the $V - I$ offset falls slightly above.

We also derived equatorial coordinates for all objects included in the data bases for the V filter. The transformation from rectangular coordinates to equatorial coordinates was derived using 79 stars identified in the USNO-A2.0 catalog. We have also compared these coordinates with those given by Magnier et al. (1992) and find a good agreement, with an average difference of 0.5 arcsec for 66 of the transformation stars found in both catalogs.

4. Selection of variables

The procedure for selecting the variables was described in detail in Paper I, so here we only give a short description, noting changes when necessary. The reduction procedure described in previous section produces databases of calibrated BVI magnitudes and their standard errors. The V database for M31F field contains 7997 stars, with up to 108 measurements, the I database contains 26540 stars with up to 69 measurements and the B database contains 3382 stars with up to 7 measurements. Figure 1 shows the distributions of stars as a function of mean $\bar{B}$, $\bar{V}$ or $\bar{I}$ magnitude. As can be seen from the shape of the histograms, our completeness starts to drop rapidly at about $\bar{B} \sim 22$, $\bar{V} \sim 22$ and $\bar{I} \sim 20.5$. The primary reason for this difference in the depth of the photometry between BV and I is the level of the combined sky and background light,

⁴IRAF is distributed by the National Optical Astronomy Observatories, which are operated by the Associations of Universities for Research in Astronomy, Inc., under cooperative agreement with the NSF




Fig. 1.— Distributions in B (dotted line), V (dashed line) and I (continuous line) of stars in the field M31F.

which is about three times higher in the I filter than in the BV filters.

The measurements flagged as “bad” (with unusually large *Daophot* errors, compared to other stars) and measurements with errors exceeding the average error, for a given star, by more than 4σ are removed. Usually zero to 10 points are removed, leaving the majority of stars with roughly $N_{good} \sim 95 - 105$ V measurements. For further analysis we use only those stars that have at least $N_{good} > N_{max}/2$ ($= 54$) measurements. There are 5838 such stars in the V database of the M31F field.

Our next goal is to select a sample of variable stars from the total sample defined above. There are many ways to proceed, and we largely follow the approach of Stetson (1996), also described in Paper I. In short, for each star we compute the Stetson’s variability index J_S (Paper I, Eq.7), and stars with values exceeding some minimum value $J_{S,min}$ are considered candidate variables. The definition of J_S is rooted in the assumption that on each visit to the program field at least one pair of observations is obtained, and only when both observations have the residual from the mean of the same sign does the pair contribute positively to the variability index. The definition of Stetson’s variability index includes the standard errors of individual observations. If, for some reason, these errors were over- or underestimated, we would either miss real variables, or select spurious variables as real ones. Using the procedure described in Paper I, we scale the *Daophot* errors to better represent the “true” photometric errors. We then select the candidate variable stars by computing the value of J_S for the stars in our V database. We used a cutoff




Fig. 2.— Variability index J_S vs. mean $\bar{V}$ magnitude for 5838 stars in the field M31F with $N_{good} > 54$. Dashed line at $J_S = 0.75$ defines the cutoff applied for variability.

of $J_{S,min} = 0.75$ and additional cuts described in Paper I to select 122 candidate variable stars (about 2% of the total number of 5838). In Figure 2 we plot the variability index J_S vs. apparent visual magnitude $\bar{V}$ for 5838 stars with $N_{good} > 54$.

5. Period determination, classification of variables

We based our candidate variables selection on the V band data collected at the MDM and the FLWO telescopes. We also have the BI -bands data for the field, up to 69 I -band epochs and up to 7 B -band epochs, although for a variety of reasons some of the candidate variable stars do not have a B or I -band counterpart. We will therefore not use the BI data for the period determination and broad classification of the variables. We will however use the BI data for the “final” classification of some variables.

Next we searched for the periodicities for all 122 candidate variables, using a variant of the Lafler-Kinman (1965) string-length technique proposed by Stetson (1996). Starting with the minimum period of 0.25 *days*, successive trial periods are chosen so

$$P_{j+1}^{-1} = P_j^{-1} - \frac{0.02}{\Delta t}, \quad (1)$$

where $\Delta t = t_N - t_1 = 398$ *days* is the time span of the series. The maximum period considered is

150 *days*. For each candidate variable 10 best trial periods are selected (Paper I) and then used in our classification scheme.

The variables we are most interested in are Cepheids and eclipsing binaries (EBs). We therefore searched our sample of variable stars for these two classes of variables. As mentioned before, for the broad classification of variables we restricted ourselves to the *V* band data. We will, however, present and use the *BI*-bands data, when available, when discussing some of the individual variable stars.

For EBs, we used the search strategy described in Paper II. Within our assumption the light curve of an EB is determined by nine parameters: the period, the zero point of the phase, the eccentricity, the longitude of periastron, the radii of the two stars relative to the binary separation, the inclination angle, the fraction of light coming from the bigger star and the uneclipsed magnitude. A total of six variables passed all of the criteria. We then went back to the CCD frames and tried to see by eye if the inferred variability is indeed there, especially in cases when the light curve is very noisy/chaotic. We decided to remove two dubious eclipsing binaries, classifying one as a periodic variable with half the determined period. Its light curve is presented in Section 6.3. The remaining four EBs with their parameters and light curves are presented in the Section 6.1.

In the search for Cepheids we followed the approach by Stetson (1996) of fitting template light curves to the data. We used the parameterization of Cepheid light curves in the *V*-band as given by Stetson (1996). There was a total of 64 variables passing all of the criteria (Paper I and II), but after investigating the CCD frames we removed 12 dubious “Cepheids”, which leaves us with 52 probable Cepheids. Their parameters and light curves are presented in Section 6.2.

After the selection of four eclipsing binaries, 52 Cepheids and one periodic variable, we were left with 65 “other” variable stars. After raising the threshold of the variability index to $J_{S,min} = 1.2$ (Paper I) we are left with 17 variables. After investigating the CCD frames we removed 10 dubious variables from the sample, which leaves seven variables which we classify as miscellaneous. Their parameters and light curves are presented in the Section 6.4.

6. Catalog of variables

In this section we present light curves and some discussion of the 64 variable stars discovered by our survey in the field M31F.⁵ The variable stars are named according to the following convention: letter V for “variable”, the number of the star in the *V* database, then the letter “D” for our project, DIRECT, followed by the name of the field, in this case (M)31F, e.g. V244 D31F.

⁵Complete *V* and (when available) *BI* photometry and 128×128 *pixel* ($\sim 40'' \times 40''$) *V* finding charts for all variables are available from the authors via the `anonymous ftp` from the Harvard-Smithsonian Center for Astrophysics and can be also accessed through the `World Wide Web`.

Tables 1, 2, 3 and 4 list the variable stars sorted broadly by four categories: eclipsing binaries, Cepheids, other periodic variables and “miscellaneous” variables, in our case meaning “variables with no clear periodicity”.

6.1. Eclipsing binaries

In Table 1 we present the parameters of the four eclipsing binaries in the M31F field. The light curves of these variables are shown in Figure 3, along with the simple eclipsing binary models discussed in the Papers I and II. The variables are sorted in the Table 1 by the increasing value of the period P . For each eclipsing binary we present its name, J2000.0 coordinates (in degrees), period P , magnitudes V_{max} , I_{max} and B_{max} of the system outside of the eclipse, and the radii of the binary components R_1 , R_2 in the units of the orbital separation. We also give the inclination angle of the binary orbit to the line of sight i and the eccentricity of the orbit e . The reader should bear in mind that the values of V_{max} , I_{max} , B_{max} , R_1 , R_2 , i and e are derived with a straightforward model of the eclipsing system, so they should be treated only as reasonable estimates of the “true” value.

One of the eclipsing binaries found, V1835 D31F, is a good DEB candidate. However, a much better light curve is necessary to accurately establish the properties of the system.

6.2. Cepheids

In Table 2 we present the parameters of 52 Cepheids in the M31F field, sorted by the period P . For each Cepheid we present its name, J2000.0 coordinates, period P , flux-weighted mean magnitudes $\langle V \rangle$ and (when available) $\langle I \rangle$ and $\langle B \rangle$, and the V -band amplitude of the variation A . In Figure 4 we show the phased B, V, I lightcurves of our Cepheids. Also shown is the best fit template lightcurve (Stetson 1996), which was fitted to the V data and then for the I data

Table 1. DIRECT ECLIPSING BINARIES IN M31F

Name (D31F)	$\alpha_{J2000.0}$ (deg)	$\delta_{J2000.0}$ (deg)	P (days)	V_{max}	I_{max}	B_{max}	R_1	R_2	i (deg)	e	Comments
V207...	10.2157	40.6399	2.5202	20.53	20.34	20.40	0.48	0.39	81	0.05	
V4320.	10.1076	40.7456	4.6694	19.46	...	19.17	0.58	0.42	70	0.00	
V1835.	10.1433	40.7184	6.3452	20.06	19.62	19.93	0.37	0.30	86	0.03	DEB
V763...	10.1807	40.7263	6.8930	19.60	19.42	19.55	0.64	0.31	63	0.02	W UMa




Fig. 3.— *BVI* lightcurves of eclipsing binaries found in the field M31F. The thin continuous line represents the best fit model for each star and photometric band. *B*-band lightcurve is shown in the bottom panel and *I*-band lightcurve (when present) is shown in the top panel.

only the zero-point offset was allowed. For the B -band data, lacking the template lightcurve parameterization (Stetson 1996), we used the V -band template, allowing for different zero-points and amplitudes. With our limited amounts of B -band data this approach produces mostly satisfactory results, but extending the template-fitting approach of Stetson (1996) to the B -band (and possibly other popular bands) would be most useful.

Some Cepheids seem to be brighter in the B band than in V . This effect is most likely caused by blending, since these variables are located in regions densely populated by stars, but could partially be due to blue binary companions of Cepheids (Evans 1994; Evans & Udalski 1994).




Fig. 4.— *BVI* lightcurves of Cepheid variables found in the field M31F. The thin continuous line represents the best fit Cepheid template for each star and photometric band. *B* (if present) is usually the faintest and *I* (if present) is usually the brightest.




Fig. 4.— Continued.




Fig. 4.— Continued.




Fig. 4.— Continued.




Fig. 4.— Continued.




Fig. 4.— Continued.




Fig. 4.— Continued.




Fig. 4.— Continued.




Fig. 4.— Continued.




Fig. 5.— BVI lightcurves of the other periodic variable found in the field M31F. B -band data (shown with the open circles) is the faintest and I is the brightest.

6.3. Other periodic variables

For one of the variables preliminarily classified as an eclipsing binary we decided upon closer examination to classify it as an “other periodic variable”. In Table 3 we present the parameters of this possible periodic variable. In Figure 5 we show its phased BVI lightcurves. We present its name, J2000.0 coordinates, period P , error-weighted mean magnitudes $\bar{V}$, $\bar{I}$ and $\bar{B}$. To quantify the amplitude of the variability, we also give the standard deviations of the measurements in the BVI bands, σ_V , σ_I and σ_B .

The period of V7438 D31F was taken to be half of the period determined by fitting a simple eclipsing binary lightcurve, so it should only be treated as a first approximation of its true value. Inspection of the V , $V - I$ and $V, B - V$ color-magnitude diagrams (Figure 6) reveals that the variable lands in the regions occupied by Cepheids. V7438 D31F has been previously identified by Baade & Swope (1965) as a Cepheid with a period of 5.12 days, very close to our value. In the P-L diagram (Figure 7), however, it is located above the region occupied by Cepheid variables, indicating that it is possibly a blend. Another fact which may favor this explanation is the small amplitude of its variability.




Fig. 6.— V , $V - I$ (upper panels) and B , $B - V$ (lower panels) color-magnitude diagrams for the variable stars found in the field M31F. The eclipsing binaries and Cepheids are plotted in the left panels and the other periodic variables and miscellaneous variables are plotted in the right panels. The dashed lines correspond to the I detection limit of $I \sim 21$ mag (upper panels) and the B detection limit of $B \sim 22.5$ mag (lower panels).

Table 2. DIRECT Cepheids in M31F

Name (D31F)	$\alpha_{J2000.0}$ (deg)	$\delta_{J2000.0}$ (deg)	P (days)	$\langle V \rangle$	$\langle I \rangle$	$\langle B \rangle$	A
V3441...	10.1201	40.7667	4.678	21.28	20.52	21.18	0.35
V4254...	10.1111	40.6766	5.718	21.48	20.50	21.99	0.36
V7832...	9.9963	40.7972	5.814	21.39	20.47	22.02	0.41
V3732...	10.1189	40.6764	6.070	20.98	19.96	21.05	0.37
V3054...	10.1243	40.7911	6.105	21.09	20.13	...	0.31
V893...	10.1716	40.7771	6.505	21.64	20.71	...	0.40
V7441...	10.0247	40.6533	6.514	21.43	19.88	...	0.37
V3860...	10.1154	40.7284	6.529	21.24	19.65	22.08	0.36
V1599...	10.1470	40.7687	6.640	20.94	19.86	21.78	0.28
V5856...	10.0799	40.6725	6.660	21.24	20.37	21.99	0.32
V5711...	10.0828	40.6802	6.707	21.07	20.10	21.87	0.43
V6623...	10.0591	40.6814	6.999	20.86	20.04	...	0.35
V5886...	10.0788	40.6903	7.458	21.12	20.54	21.91	0.36
V6406...	10.0662	40.6569	7.563	20.95	20.24	...	0.35
V3289...	10.1232	40.7453	7.599	20.89	19.76	20.77	0.27
V5893...	10.0797	40.6513	7.655	21.00	20.02	21.70	0.34
V6962...	10.0462	40.6923	7.782	21.31	20.47	...	0.24
V7741...	10.0076	40.6885	8.099	21.03	20.17	20.81	0.24
V6098...	10.0748	40.6395	8.471	20.52	19.52	...	0.28
V4855...	10.0966	40.8035	9.064	20.70	19.70	...	0.30
V5498...	10.0883	40.6611	9.387	20.72	20.27	20.98	0.24
V7074...	10.0388	40.7778	9.478	20.78	19.70	21.51	0.30
V5097...	10.0967	40.6804	9.662	20.66	19.92	21.16	0.25
V6483...	10.0637	40.6739	9.736	20.83	20.13	21.20	0.23
V5994...	10.0774	40.6532	9.886	20.67	19.53	21.45	0.22
V4556...	10.1054	40.7022	9.894	20.54	19.41	21.20	0.25
V6195...	10.0722	40.6463	9.924	21.21	20.05	22.54	0.35
V5178...	10.0949	40.6807	9.932	20.30	19.30	20.18	0.22
V7393...	10.0278	40.6364	9.937	20.50	19.51	21.15	0.31
V3550...	10.1188	40.7607	10.468	20.55	19.85	21.03	0.32
V2320...	10.1341	40.7656	10.868	20.27	19.54	20.60	0.34
V4125...	10.1109	40.7332	11.139	20.46	19.63	21.11	0.31
V1549...	10.1488	40.7569	11.764	20.72	19.68	21.49	0.35
V5442...	10.0892	40.6766	11.902	19.65	18.58	19.45	0.16
V5696...	10.0806	40.7520	12.287	20.74	19.54	21.62	0.43
V5598...	10.0855	40.6741	12.311	20.34	19.45	20.96	0.37
V6267...	10.0684	40.6995	12.324	20.38	19.30	19.27	0.17
V2156...	10.1353	40.7925	12.831	20.37	19.40	20.76	0.27
V3373...	10.1199	40.7971	12.872	20.94	19.55	21.81	0.41
V4955...	10.0959	40.7808	13.034	20.70	19.67	21.45	0.41
V1633...	10.1461	40.7760	13.043	21.30	19.85	...	0.45




Fig. 7.— Diagram of $\log P$ vs. I for the Cepheids (open circles), RV Tau (dotted circles) variables and the other periodic variable (filled circle). The sizes of the circles are proportional to the V amplitude of the variability.

6.4. Miscellaneous variables

In Table 4 we present the parameters of seven miscellaneous variables in the M31F field, sorted by increasing value of the mean magnitude $\bar{V}$. In Figure 8 we show the unphased VI lightcurves of the miscellaneous variables. For each variable we present its name, J2000.0 coordinates and mean magnitudes $\bar{V}$, $\bar{I}$ and $\bar{B}$. To quantify the amplitude of the variability, we also give the standard deviations of the measurements in BVI bands, σ_V , σ_I and σ_B . In the “Comments” column we give a rather broad sub-classification of the variability.

All of the variables seem to represent the LP type of variability. A closer inspection of the color-magnitude diagrams (Figure 6) reveals that three variables (V667, V1229 and V2285 D31F) land in the same area as Cepheids. Based on their lightcurves it was possible to roughly estimate the periods of the first two to be around 90 and 100 days, respectively. Using these periods to place the stars on the P-L diagram (Figure 7) suggests they may be RV Tauri type variables. V1665 and V1724 D31F are most likely Mira-type variables, based on their location in the color-magnitude diagrams.

6.5. Comparison with other catalogs

The area of the M31F field coincides with two overlapping fields observed by Baade. The catalogs of variable stars discovered in those fields are given by Gaposchkin (1962, field II) and Baade & Swope (1965, field III). We succeeded in the cross-identification of all but one of the 55 Cepheid variables found in field III with stars on our template. We have discovered independently 24 of those Cepheids and found a very good agreement between the period determinations. We have also confirmed the periods of 23 other Cepheids, which eluded our detection, in large part due to their faintness and the strict criteria we have imposed in our process of Cepheid selection (see Table 5 for cross-ids).

Out of the 38 unique Cepheid variables listed in the field II catalog, located within our M31F field, we have found 14 Cepheids and confirmed the periods of two. The remaining field II Cepheids have evaded positive cross-identification with our template stars.

Another overlapping variable star catalog is given by Magnier et al. (1997, hereafter Ma97). Out of the three variable stars in Ma97 which are in our M31F field, we cross-identified one, also classifying it as a Cepheid. The other two did not qualify as variable star candidates because of low J_S values.

Table 2—Continued

Name (D31F)	$\alpha_{J2000.0}$ (deg)	$\delta_{J2000.0}$ (deg)	P (<i>days</i>)	$\langle V \rangle$	$\langle I \rangle$	$\langle B \rangle$	A
V1619...	10.1503	40.6545	13.141	20.61	19.45	21.47	0.34
V4682...	10.1046	40.6552	13.297	20.73	19.30	21.72	0.34
V6640...	10.0593	40.6558	13.761	19.67	...	...	0.18
V4861...	10.0972	40.7849	13.991	20.47	19.31	21.22	0.41
V6503...	10.0615	40.7215	15.210	20.77	19.55	21.44	0.38
V4708...	10.1039	40.6631	15.690	20.50	19.33	21.38	0.35
V1893...	10.1415	40.7417	16.756	18.79	17.47	19.54	0.18
V6208...	10.0714	40.6561	17.572	19.74	18.78	20.28	0.48
V821....	10.1760	40.7640	20.547	21.35	19.61	...	0.55
V602....	10.1880	40.7392	31.416	21.05	19.12	...	0.31
V2203...	10.1369	40.7306	55.373	17.94	17.17	18.32	0.17

Table 3. DIRECT Other Periodic Variables in M31F

Name (D31F)	$\alpha_{J2000.0}$ (deg)	$\delta_{J2000.0}$ (deg)	P (<i>days</i>)	$\bar{V}$	$\bar{I}$	$\bar{B}$	σ_V	σ_I	σ_B	Comments
V7438...	10.0252	40.6442	5.1	20.41	19.41	21.02	0.15	0.11	0.13	Cepheid?

Note. — V7438 was identified by Baade & Swope (1965) as Cepheid variable 232 with $P = 5.12$ days.




Fig. 8.— *VI* lightcurves of the miscellaneous variables found in the field M31F. *I* (if present) is plotted in the two right panels. *B*-band data is not shown.

Table 4. DIRECT Miscellaneous Variables in M31F

Name (D31F)	$\alpha_{J2000.0}$ (deg)	$\delta_{J2000.0}$ (deg)	$\bar{V}$	$\bar{I}$	$\bar{B}$	σ_V	σ_I	σ_B	Comments
V244...	10.2103	40.7380	18.39	16.68	19.87	0.12	0.07	0.05	LP
V1665.	10.1460	40.7562	19.27	16.01	21.40	0.20	0.16	0.10	LP
V1724.	10.1446	40.7499	19.63	16.34	0.00	0.24	0.13	0.00	LP
V2285.	10.1373	40.6841	19.67	18.81	19.94	0.08	0.07	0.08	RV Tau?
V1229.	10.1590	40.7389	20.47	19.57	21.07	0.23	0.10	0.18	RV Tau?
V764...	10.1775	40.8110	20.51	0.00	0.00	0.35	0.00	0.00	LP
V667...	10.1852	40.7265	21.18	19.33	0.00	0.35	0.22	0.00	RV Tau?




Fig. 8.— Continued.

Table 5. Cross-Identifications of the DIRECT Cepheid Variables in M31F

Name (D31F)	P (<i>days</i>)	field II	P (<i>days</i>)	field III	P (<i>days</i>)	Other	P (<i>days</i>)
V3441...	4.678			108	5.000		
V4254...	5.718	351	5.719	153	5.721		
V3732...	6.070	352	6.071	175	6.074		
V3054...	6.105			92	6.101		
V7441...	6.514	230	6.508				
V3860...	6.529			24	6.526		
V1599...	6.640			145	6.637		
V5856...	6.660	328	6.660	30	6.700		
V5711...	6.707	330	6.709	29	6.708		
V6623...	6.999	326	7.000				
V5886...	7.458	332	7.457	70	7.463		
V6406...	7.563	319	7.553				
V5893...	7.655	320	7.823				
V6962...	7.782	225	7.780				
V7741...	8.099	222	8.094				
V6098...	8.471	315	8.464				
V4855...	9.064			51	9.085		
V5097...	9.662	348	9.662	68	9.678		
V6483...	9.736	325	9.483				
V4556...	9.894	341	9.881	60	9.870		
V6195...	9.924	318	9.921				
V7393...	9.937	234	9.933			Ma97 4	9.0
V3550...	10.468			72	10.461		
V2320...	10.868			109	10.858		
V4125...	11.139			20	11.147		
V1549...	11.764			54	11.766		
V5696...	12.287			17	12.286		
V5598...	12.311	329	12.312	135	12.358		
V6267...	12.324	334	12.294	133	12.284		
V2156...	12.831			128	12.821		
V4955...	13.034			14	13.051		
V1633...	13.043			107	13.021		
V1619...	13.141	415	13.125	31	0.000		
V4682...	13.297	357	13.293				
V4861...	13.991			74	13.966		
V6503...	15.210	339	15.232	150	15.216		
V4708...	15.690	355	15.699	114	15.625		
V6208...	17.572		17.569			H22	17.60

Note. — field II refers to the catalog published by Gaposchkin (1962); field III - Baade & Swope (1965); Ma97 - Magnier et al. (1997); H - Hubble(1929)




Fig. 9.— Location of eclipsing binaries (filled squares) and Cepheids (open circles) in the field M31F, along with the blue stars ($B - V < 0.4$) selected from the photometric survey of M31 by Magnier et al. (1992) and Haiman et al. (1994). The sizes of the circles representing the Cepheids variables are proportional to the logarithm of their period. Fields II and III observed by Baade are marked with dashed lines.

7. Discussion

In Figure 6 we show V , $V - I$ and V , $B - V$ color-magnitude diagrams for the variable stars found in the field M31F. The eclipsing binaries and Cepheids are plotted in the left panels and the other periodic variables and miscellaneous variables are plotted in the right panels. As expected, the eclipsing binaries occupy the blue upper main sequence of M31 stars. The Cepheid variables group near $B - V \sim 1.0$, with considerable scatter probably due to the differential reddening across the field. The other periodic variable is located on the CMD in the part occupied by Cepheids. The miscellaneous variables are scattered throughout the CMDs and represent several classes of variability. Two of them are very red with $V - I > 2.0$, and are probably Mira variables.

In Figure 9 we plot the location of eclipsing binaries and Cepheids in the field M31F, along with the blue stars ($B - V < 0.4$) selected from the photometric survey of M31 by Magnier et al. (1992) and Haiman et al. (1994). The sizes of the circles representing the Cepheids variables are proportional to the logarithm of their period. As could have been expected, both types of variables group along the spiral arms, as they represent relatively young populations of stars. Many Cepheid variables are located in the star-forming region NGC206. We will explore various properties of our sample of Cepheids in the future paper (Sasselov et al. 1999, in preparation).

We would like to thank the TAC of the F. L. Whipple Observatory (FLWO) and the TAC of the Michigan-Dartmouth-MIT (MDM) Observatory for the generous amounts of telescope time devoted to this project. We are grateful to Bohdan Paczyński for motivating us to undertake this project and always helpful comments and suggestions. We thank Lucas Macri for taking some of the data described in this paper, Przemek Woźniak for his FITS-manipulation programs and Eugene Magnier for the Cepheid catalogs. The staff of the MDM and the FLWO observatories is thanked for their support during the long observing runs. KZS was supported by the Harvard-Smithsonian Center for Astrophysics Fellowship. JK was supported by NSF grant AST-9528096 to Bohdan Paczyński and by the Polish KBN grant 2P03D011.12.

REFERENCES

- Andersen, J. 1991, *A&AR*, 3, 91
- Baade, W., & Swope, H. H. 1963, *AJ*, 68, 435
- Baade, W., & Swope, H. H. 1965, *AJ*, 70, 212
- Crotts, A. P. S., & Tomaney, A. B. 1996, *ApJ*, 473, L87
- Evans, N. R., & Udalski, A. 1994, *AJ*, 108, 653
- Evans, N. R. 1994, *ApJ*, 436, 273
- Freedman, W. L., & Madore, B. F. 1990, *ApJ*, 365, 186
- Freedman, W. L., Wilson, C. D., & Madore, B. F. 1991, *ApJ*, 372, 455
- Gaposchkin, S. 1962, *AJ*, 67, 334
- Guinan, E. F., et al. 1998, *ApJ*, 509, L21
- Haiman, Z., et al. 1994, *A&A*, 286, 725
- Hilditch, R. W. 1996, in: *ASP Conf. Ser. 90, The Origins, Evolution and Destinies of Binary Stars in Clusters*, ed. E. F. Milone & J.-C. Mermilliod (San Francisco: ASP), 207
- Holland, S. 1998, *AJ*, 115, 1916
- Hubble, E. 1926, *ApJ*, 63, 236
- Hubble, E. 1929, *ApJ*, 69, 103
- Huterer, D., Sasselov, D. D., Schechter, P. L. 1995, *AJ*, 110, 2705
- Jacoby, G. H., et al. 1992, *PASP*, 104, 599
- Kaluzny, J., Stanek, K. Z., Krokenberger, M., Sasselov, D. D., Tonry, J. L., & Mateo, M. 1998, *AJ*, 115, 1016 (Paper I)
- Kaluzny, J., Mochejska, B. J., Stanek, K. Z., Krokenberger, M., Sasselov, D. D., Tonry, J. L., & Mateo, M. 1999, *AJ*, in press (Paper IV) (astro-ph/9902382)

- Krockenberger, M., Sasselov, D. D., & Noyes, R. 1997, *ApJ*, 479, 875
- Lafler, J., & Kinman, T. D. 1965, *ApJS*, 11, 216
- Landolt, A. 1992, *AJ*, 104, 340
- Magnier, E. A., Augusteijn, T., Prins, S., van Paradijs, J., & Lewin, W. H. G. 1997, *A&AS*, 126, 401 (Ma97)
- Magnier, E. A., Lewin, W. H. G., Van Paradijs, J., Hasinger, G., Jain, A., Pietsch, W., & Truemper, J. 1992, *A&AS*, 96, 37
- Metzger, M. R., Tonry, J. L., & Luppino, G. A. 1993, in *ASP Conf. Ser. 52, Astronomical Data Analysis Software and Systems II*, ed. R. J. Hanisch, R. J. V. Brissenden, & J. Barnes, (San Francisco: ASP), 300
- Monet, D., et al. 1996, *USNO-SA2.0*, (U.S. Naval Observatory, Washington DC).
- Paczyński, B. 1997, in *The Extragalactic Distance Scale*, ed. M. Livio, M. Donahue & N. Panagia (Cambridge: Cambridge Univ. Press), 273
- Sasselov, D. D., et al. 1999, in preparation
- Stanek, K. Z., Kaluzny, J., Krockenberger, M., Sasselov, D. D., Tonry, J. L., & Mateo, M. 1998, *AJ*, 115, 1894 (Paper II)
- Stanek, K. Z., Kaluzny, J., Krockenberger, M., Sasselov, D. D., Tonry, J. L., & Mateo, M. 1999, *AJ*, in press (Paper III) (astro-ph/9901331)
- Stanek, K. Z., & Garnavich, P. M. 1998, *ApJ*, 503, L131
- Stetson, P. B. 1987, *PASP*, 99 191
- Stetson, P. B. 1992, in *ASP Conf. Ser. 25, Astrophysical Data Analysis Software and Systems I*, ed. D. M. Worrall, C. Bimesderfer, & J. Barnes (San Francisco: ASP), 297
- Stetson, P. B. 1996, *PASP*, 108, 851
- Szentgyorgyi, A., et al. 1999, in preparation
- Tonry, J. L., Blakeslee, J. P., Ajhar, E. A., & Dressler, A., 1997, *ApJ*, 475, 399
- Udalski, A., Pietrzyński, G., Woźniak, P. R., Szymański, M., Kubiak, M., & Żebruń, K., 1998, *ApJ*, 509, L25